

Original Research Article

## Effect of Pranayama on Body Mass Index in Young Medical Students

Patil Smita V<sup>1</sup>, Gaikwad Pandurang B<sup>2</sup>, Suvarna T. Jadhav<sup>3</sup>, Nilima R. Patil<sup>4</sup>, Vinayak B. Gaikwad<sup>5</sup>

<sup>1</sup>Assistant Professor, Dept. of Physiology, Bharati Vidyapeeth University Dental College & Hospital, Sangli.

<sup>2</sup>Assistant Professor, Dept. of Physiology, Prakash Institute of Medical Science & Research Islampur, Dist. Sangli (MS).

<sup>3</sup>Assistant Professor, Dept of Biochemistry, Bharati Vidyapeeth University Dental College & Hospital, Sangli.

<sup>4</sup>Assistant Professor, Dept. of Microbiology, Bharati Vidyapeeth University Dental College & Hospital, Sangli.

<sup>5</sup>Statistician Cum Tutor, Dept. of Community Medicine, B.K.L Walawalkar, Rural Medical Collage.

Corresponding Author: Patil Smita V.

Received: 29/01/2016

Revised: 05/02/2016

Accepted: 15/02/2016

### ABSTRACT

**Background:** Association between Body Mass Index (BMI) and cardiovascular diseases has been consistently observed, but yet it is not clearly understood, possibly because of interactions with other influencing factors. One unresolved question is whether there is a linear relationship.

**Aim:** The aim of the study was to investigate the effect of pranayama on BMI in young medical students.

**Materials & methods:** A comparative cross sectional study was conducted on 60 young volunteers (42- male, 18- female) of age group 17-20years, to study the effect of pranayama on BMI. Anthropometric parameters like height (cm), weight (Kg) were recorded. BMI was calculated using formula,  $BMI = Wt \text{ (kilograms)}/Ht^2 \text{ (meters)}$  before and after 4<sup>th</sup>, 8<sup>th</sup>, & 12<sup>th</sup> week of pranayama practice. ANOVA and Post hoc Tukey's test was applied for statistical analysis and p-value <0.05 was considered the level of significance.

**Results:** Mean BMI Before pranayama practice was  $21.657 \pm 3.01$ . There was significant decrease in BMI ( $20.998 \pm 2.76$ ) after 12<sup>th</sup> week of pranayama practice. There was no significant change in BMI observed after 4<sup>th</sup> week of pranayama practice But significant decreased in BMI occur after 8<sup>th</sup> & 12<sup>th</sup> week of pranayama.

**Conclusion:** Short term pranayama have reducing impact on BMI and also have positive and useful effect on weight reduction in young adults.

**Key words:** Pranayama, Height, Weight, Body Mass Index.

### INTRODUCTION

Over thousands of years asanas, pranayama, and meditation are three main techniques of yoga practiced in India, to attain functional harmony between the body and mind. With increased awareness and interest in health and natural remedies, yogic techniques including pranayama are getting importance and becoming increasingly acceptable to the scientific society. <sup>[1]</sup> Obesity is the newer chronic non-

communicable disease, though is increasing in an epidemic proportion is one of the today's most neglected public health problems according to the World Health Organization. <sup>[2]</sup> A number of clinical measurements for obesity have been used to determine susceptibility to different diseases. <sup>[3]</sup> These include anthropometric indices such as body mass index (BMI), waist-hip ratio (WHR) and waist circumference (WC). Body Mass Index

(BMI) is considered to be one of the most useful indices for obesity in adults. BMI is determined by dividing weight (wt) in kilograms by height (ht) in meters square.

Nowadays Obesity is considered as global epidemic. It cause a large number of health problems. Other than the genetic predisposition, favorable environment for development of obesity due to sedentary life style, inappropriate intake of caloric rich easily available junk food and automated working profile. Various measures with variable results are used to reduce weight like dieting, hydrotherapy, steam bath, pharmacological therapy, surgical therapy, etc.

Hence present study was conducted to find out the effect of short term pranayama on weight reduction & body mass index in young medical students which will scientifically contribute to identify at-risk population well in advance and will also help to implement necessary action to obtain desired physical fitness in the form of optimum body composition and thereby to prevent / delay future health hazards in young adults.

## MATERIALS AND METHODS

A cross sectional study was carried out on healthy 1st year M.B.B.S. students in Dept. of Physiology G.M.C. Miraj (Maharashtra). Students were evaluated as per standard proforma, which included a questionnaire. The students with, past or present history suggestive of cardiovascular or respiratory illness or any other systemic illness, any family history of asthma or allergic diseases, were not included in the study. Only nonsmoker students were enrolled. Only 60 students were eligible to whom the experimental protocol was

explained and written informed consent was obtained from them. Out of sixty students 42 were male and 18 were females.

In the beginning itself the institutional ethical committee approval was obtained for the study. Anthropometric measurements: Height (in meters) - using a stadiometer, weight (in kilograms) - using a digital weighing scale (precision of 100 grams), were measured. Body mass index (BMI) was calculated using the formula,

$$BMI = Wt \text{ (kilograms)} / Ht^2 \text{ (meters)}$$

After recording the above parameters, students were trained by yoga instructor. They Performed the Pranayama practice daily in the evening for one hour (5.15 pm - 6.15 pm), Six days in a week for three months under experts' supervision. The Pranayama practice consisted of prayer, Nadishuddhi, Anulom-vilom, Savitri, Sitakari, Sadanta, shawasan Bhastrika, Omkar recitation, each done for a period of 5 min. followed by Bhramari, Kapalhati for a period of 10 min.

All the parameters were recorded After 4 weeks, 8 weeks and 12 weeks of Pranayama practice & Results were presented as Mean SD. Repeated measure ANOVA test was used to find the significance of study parameters by using SPSS 16.0 version. P value less than 0.05 was considered statistically significant.

## RESULTS

Table No- I Percentage distribution of students according to BMI

Sr. No.	BMI (kg/m <sup>2</sup> ) category	Prevalence (%)
1	Under weight BMI < 18.5	8.3
2	Normal weight BMI 18.5-24.9	83.3
3	Over weight BMI 25 to <30	5
4	Obese BMI 30 & above	3.3

Table No- II Comparison of Mean ± SD Values Height and weight before and after 4<sup>th</sup>, 8<sup>th</sup> & 12<sup>th</sup> week of Pranayama practice using Repeated measure ANOVA.

Parameters	Before pranayama Mean ± SD	After 4 <sup>th</sup> week of pranayama Mean ± SD	After 8 <sup>th</sup> week of pranayama Mean ± SD	After 12 <sup>th</sup> week of pranayama Mean ± SD
Height (cm)	165.30±7.046	165.30±7.046	165.30±7.046	165.30±7.046
Weight (Kg)	59.25±9.502	59.07±9.331	58.46±9.401	58.26±9.341

**Table No.-III Comparison of Mean  $\pm$  SD Values of BMI before and after 4<sup>th</sup>, 8<sup>th</sup> & 12<sup>th</sup> week of Pranayama practice using Repeated measure ANOVA.**

Parameters	BMI(kg/m <sup>2</sup> ) Mean $\pm$ SD	F Value	'P' Value
Before pranayama	21.657 $\pm$ 3.01	16.36	0.001*
4 <sup>th</sup> week of after pranayama	21.597 $\pm$ 2.98		
8 <sup>th</sup> week of after pranayama	21.383 $\pm$ 3.07		
12 <sup>th</sup> week of after pranayama	20.998 $\pm$ 2.76		

**Table No.-IV Between group Comparison of BMI before and with 4<sup>th</sup>, 8<sup>th</sup> & 12<sup>th</sup> week of after Pranayama practice using Post hoc tukeys' test.**

Parameters	After 4 <sup>th</sup> week of pranayama		After 8 <sup>th</sup> week of pranayama		After 12 <sup>th</sup> week of pranayama	
	Mean difference	'P' Value	Mean difference	'P' Value	Mean difference	'P' Value
BMI	0.060	0.319 (NS)	.274	0.006*	0.659	0.001*

\*-significant, NS- Not significant

## DISCUSSION

The pattern of body fat distribution is recognized as an important predictor of the health risks of obesity. Individuals with more fat on the trunk, especially abdominal fat, are at increased risk of obesity related health problems compared with individuals who are equally fat, but have more of their fat on extremities.

In present study, we observed that body weight & body mass index decreased significantly. Decrease in body weight causes change in body fat distribution. [4,5] Mauro Zamboni *et al* reported that weight loss is associated with changes in regional fat distribution. [6]

In the present study as shown in table no. II before pranayama practice BMI was 21.657(kg/m<sup>2</sup>) and, when compared BMI after 4<sup>th</sup>, 8<sup>th</sup>, & 12<sup>th</sup> week of pranayama practice it was 21.597(kg/m<sup>2</sup>), 21.383 (kg/m<sup>2</sup>) & 20.998 (kg/m<sup>2</sup>) respectively. In this study we found that after 8<sup>th</sup> & 12<sup>th</sup> week pranayama practice BMI was significantly decreased (P=0.001). But there was no statistically significant decrease in BMI was observed after 4<sup>th</sup> week of pranayama practice.

The results of this study are coincide with Manchanda *et al.*, (2000), [7] they had observed weight reduction-6.8 $\pm$ 8.2% (P=0.0019) after yogic lifestyle intervention in coronary atherosclerotic patients. Similarly Schmidt *et al*; (1997) [8] studied cardiovascular risk factors and hormones during comprehensive residential 3 months kriya yoga training and they observed that significant reduction in body mass index.

Calle-Pascual *et al.* (1991) [9] studied behaviour modification in obese subjects with type-2 diabetes mellitus and observed that BMI reduced from 34.2 $\pm$ 0.8 kg/m<sup>2</sup> to 30.6 $\pm$ 1.1 kg/ m<sup>2</sup> (P=0.05).

Our study also coincide with various other studies, they have shown that effect of pranayama on obesity in the form of weight reduction. Nirmala N. Nayak *et al* (2004) [10] studied that various yoga asanas including Kapalbhathi seem to have a positive effect in reducing obesity. Swami Ramdev mentioned that Kapalbhathi is helpful in reducing obesity (2005). [11] Dinkar R. Kekan (2013) reported that Kapalbhathi has reducing impact on Body mass index and abdominal skinfold thickness in overweight individuals. [12]

The possible reason for the reduction of weight and BMI in present study may be due to pranayama involves abdominal muscle contractions with forceful exhalation and natural inhalation. It is a form of abdomino-respiratory-autonomic exercise. Due to this, respiratory, abdominal and gastrointestinal receptors get stimulated. Also afferents, centres in brainstem, cortex and effectors get stimulated. This leads to synchronous stimulation of autonomic nervous system, hypothalamus, pineal gland and other associated brain structures increases synchronous discharge to all parts of the body including endocrine and metabolic processes. This is responsible for the effect of pranayama on fat metabolism. This causes increase in basal metabolic rate, and because of this there is increase in calories consumption and decrease in fat deposition and so reduction in weight. This

might be the possible reason behind reduction in BMI in present study.

## CONCLUSION

It is concluded that the short term pranayama have positive and useful effect on certain cardiovascular risk factors like obesity and weight reduction.

## REFERENCES

1. N. K. Subbalakshmi, S. K. Saxena. - Urmimala, an Urban J.A. D'Souza. Immediate effect of Nadishodhan pranayama on some selected parameters of cardiovascular, pulmonary, and higher functions of brain. *Thai Journal of Physiological Sciences*. Vol. 18(2): 10-16, 2005.
2. Pednekar MS. Association of body mass index with all-cause and cause-specific mortality: Findings from a prospective cohort study in Mumbai (Bombay), India. *Int J Epidemiol*: 2008; 37: 524-35.
3. Cameron, A.J., Welborn, T.A. and Zimmet P.Z. (2003). Overweight and obesity in Australia: The 1999-2000 Australian Diabetes, Obesity and Lifestyle Study. *Medical Journal of Australia* 178, pp.427-432.
4. Francis E. Johnston, Thomas A. Wadden, Albert J. Stunkard, Manual Pefla, Jack Wang, Richard N. Pierson, and Theodore B. Van Itallie. - Body fat deposition in adult obese women. I Patterns of fat distribution 1-3. *Am J Clin Nutr*. 47:225-228, 1988.
5. Thomas A. Wadden, Albert J. Stunkard, Francis E. Johnston, Jack Wang, et al. - Body fat deposition in adult obese women. II Changes in fat distribution accompanying weight reduction 1-3. *Am J Clin Nutr*. 47:229-234, 1988.
6. Mauro Zamboni, Fabio Armellini, Emanuela Turcato et al. - Effect of weight loss on regional body fat distribution in premenopausal women 1, 2. *Am J Clin Nutr*. 58:29-34, 1993.
7. Manchanda SC, Narang R, Reddy KS, Sachdeva U, Prabhakaran D, Dharmananda S, Rajani M, Bijalani RL(2000). Coronary Atherosclerotic reversal potential of yoga life style intervention *JAPI* 48(7)687-689.
8. Schmidt T, Wijga A, Von ZurMuhlen A, Brabant G, Wagner TO (1997). Changes in cardiovascular risk factor and hormones during a comprehensive residential three month kriya yoga training and vegetarian nutrition. *Acta Physiol Scand Suppl* 640 158-162. Tiwari OP (1983). Yoga for keeping fit in old age, Swastha Hind.
9. Calle-Pascual AL, Rodriguez C, Camacho F, Sanchez R, Martin-Alvarez E Yuste, Hidalgo I, Diaz RJ, Calle JR, Charro AL(1992). Behaviour modification in obese subjects with type 2 diabetes mellitus. *Diabetes Research and clinical practice* 15 157-162.
10. Nirmala N. Nayak, Kamala Shankar. - Yoga: a therapeutic approach. *Phys Med Rehabil Clin N Am*. 15: 783 - 798, 2004.
11. Swami Ramdev. - Seven Kinds of Pranayama. In: *Pranayama - Its Philosophy and Practice*. Divya Prakashan, Haridwar. 28-29, May 2005.
12. Dinkar R. Kekan Effect of Kapalbhathi Pranayama on Body Mass Index and Abdominal Skinfold Thickness *Indian Medical Gazette* 2013, 424- 427.

How to cite this article: Patil SV, Gaikwad PB, Jadhav ST et al. Effect of pranayama on body mass index in young medical students. *Int J Res Rev*. 2016; 3(2):24-27.

\*\*\*\*\*