

Nausea and Vomiting in Pregnancy

Dr. Neerja Sharma

Specialist Gynaecology, South Delhi MCD

ABSTRACT

Nausea and vomiting in pregnancy also known as “Morning Sickness” usually begins between the 4th and 7th weeks in 80% of pregnant women and resolves by the 20th week of gestation. It is usually a mild, self limited condition which is often controlled by conservative measures. The etiology of nausea and vomiting in pregnancy remains unknown. There are different factors and theories believed to cause this. Management includes non-pharmacological (e.g. assurance/counselling, acupressure, dietary measures and music therapy) and pharmacological therapy (like medication, antiemetics, antihistamines & anticholinergics, motility drugs and corticosteroids).

Key words: Nausea, Vomiting, Pregnancy, Morning sickness.

INTRODUCTION

Nausea and vomiting in pregnancy are also known as “Morning Sickness”. Morning Sickness is nausea that occurs during pregnancy. The name is a misnomer as it can strike at any time of the day or night. It usually begins between the 4th and 7th weeks in 80% of pregnant women and resolves by the 20th week of gestation. It is usually a mild, self limited condition which is often controlled by conservative measures. Rarely morning sickness is so severe that it is classified as Hyperemesis Gravidarum. Hyperemesis Gravidarum includes persistent vomiting dehydration, ketosis, electrolyte disturbances and weight loss.

CAUSES

The etiology of nausea and vomiting in pregnancy remains unknown. Many believe that psychologic factors are responsible for nausea and vomiting in pregnancy but few data support this theory. Many reports have suggested that hormones may cause nausea and vomiting. Elevated levels of Human Chorionic Gonadotropin

(HCG) and suppressed levels of Thyrotropin stimulating hormones were found in patients with Hyperemesis Gravidarum. A recent study [1] has suggested that chronic infection with Helicobacter Pylori may play a role in Hyperemesis Gravidarum.

Outcome

Women with morning sickness have been noted to have improved pregnancy outcomes.

Treatment

Non-pharmacologic therapy

Assurance - Although morning sickness is not strongly associated with psychologic illness, appropriate support from family members as well as from medical staff is recommended.

Acupressure [3,4] - Acupressure is an intervention without known adverse effects. The most common point is (P6) Pericardium 6 or Nei guan Point, which is located three finger breadths above the wrist of the volar surface.

Dietary measures- Maximum pregnant women are relieved by simple dietary

modifications. Intake of frequent, small meals and to avoid fatty, oily food helps the women with morning sickness. Solid foods should be bland tasting, high in

carbohydrates. Ginger [5] is a popular alternative treatment for morning sickness. Music therapy- when used with standard treatment, music therapy can help to reduce morning sickness

Differential Diagnosis and Evaluation (Table 1 [2])

Gastrointestinal disorders	Metabolic disorders
• Gastroenteritis	• Diabetic ketoacidosis
• Biliary tract disease	• Porphyria
• Hepatitis	• Addison's disease
• Intestinal obstruction	• Hyperthyroidism
• Peptic ulcer disease	• Neurologic disorders
• Pancreatitis	• Pseudotumor cerebri
• Appendicitis	• Vestibular lesions
Genitourinary tract disorders	• Migraine headaches
• Pyelonephritis	• Central nervous system tumors
• Uremia	• Pregnancy-related conditions
• Degenerating uterine leiomyoma	• Nausea and vomiting in pregnancy*
• Torsion	• Acute fatty liver in pregnancy
• Kidney stones	• Pre-eclampsia
	• Drug toxicity or intolerance

* including Hyperemesis Gravidarum

Pharmacologic Therapy

Pyridoxine (Vitamin B₆) – Pyridoxine alone or in combination with Doxylamine 1 Tab 6-8 hourly very effective in relieving the symptoms. Pyridoxine-

Doxylamine is still the only medication that the US Food and Drug Administration has specifically labeled for the treatment of nausea and vomiting in pregnancy.

Medication	Dosages*	Pregnancy category
• Pyridoxine (vitamin B ₆)	25 mg orally three times daily	A #
• Doxylamine (unisom)	25 mg orally once daily	\$
Antiemetics		
• Chlorpromazine	10-25 mg orally 2-4 times daily	C
• Prochlorperazine	5-10 mg orally 3 or 4 times daily	C
• Promethazine	12.5-25 mg orally every 4-6 hours	C
• Trimethobenzamide	250 mg orally 3 or 4 times daily	C
• Ondansetron	8 mg orally 2 or 3 times daily	B
• Droperidol	0.5-2 mg IV or IM every 3 or 4 hours	C
Antihistamines and anticholinergics		
• Diphenhydramine	25-50 mg orally every 4-8 hours	B
• Meclizine	25 mg orally every 4-6 hours	B
• Dimenhydrinate	50-100 mg orally every 4-6 hours	B
Motility drug		
• Metoclopramide	5-10 mg orally every 3 times daily	B
Corticosteroid		
• Methylprednisolone	16 mg orally 3 times daily; then taper	C


Abbreviations: IV, intravenously; IM, intramuscularly

*these regimens usually are administered only as needed

#The pregnancy category for doxylamine relates to its use as a vitamin supplement.

\$ According to the physicians' desk reference for nonprescription drugs and dietary supplements, doxylamine should not be taken by pregnant women or women who are nursing a baby; however, some research supports its efficacy and safety.

Flow Chart 1: An approach to the management of women with nausea and vomiting in pregnancy



REFERENCES

- Hayakawa S, Nakajima N, Karasaki-Suzuki M, Yoshinaga H, Arakawa Y, Satoh K, et al. Frequent presence of *Helicobacter pylori* genome in the saliva of patients with hyperemesis gravidarum. *Am J Perinatol*, 2000; 17(5):243-7.
- Goodwin TM. Hyperemesis gravidarum. *Clin Obstet Gynecol*. 1998;41(3):597-605.
- Hyde E. Acupressure therapy for morning sickness. A controlled clinical trial. *J Nurse Midwifery*. 1989;34(4):171-8.
- de Aloysio D, Penacchioni P. Morning sickness control in early pregnancy by Neiguan point acupressure. *Obstet Gynecol*. 1992; 80(5):852-4.
- Fischer-Rasmussen W, Kjaer SK, Dahl C, Asping U. Ginger treatment of hyperemesis gravidarum. *Eur J Obster Gynecol Reprod Biol*. 1991;38(1):19-24.

6. Green SM (Ed). Tarascon pocket pharmacopoeia. Loma Linda: Tarascon;2002.
7. Physicians' desk reference for nonprescription drugs and dietary supplements, 22nd ed. Montvale: Medical Economics; 2001.

How to cite this article: Sharma N. Nausea and vomiting in pregnancy. International Journal of Research and Review. 2017; 4(4):48-51.

